How to rectify scanned air photos using ERDAS Imagine AutoSync (in the real world)

Carrie Guiles
Spatial Analysis Center
Yellowstone NP

Problem: road is being attacked by streams



Problem: road is being attacked by streams



What imagery do we have?

- 2009 NAIP
- 2006 NAIP
- 2006 QuickBird
- 2002 CIR DOQQ
- 1996 Grayscale DOQQ
- And...

...a lot of paper photos













So here's the plan:

- Scan air photos
- ■Rectify them ←
- Extract stream courses
- See how the stream changed over time

Options for rectification?

- No camera reports for most
- Multiple scales
- Not a lot of time
- Have an AutoSync license, why not use it?

AutoSync looks so easy

- But it really isn't
- This process, like so much of image processing is a bit of science and a lot of art.
- The ERDAS documentation for this module will help you understand the process - I would suggest scanning over the following - C:\ERDAS\ERDAS Desktop 2010\help\hardcopy\ AutoSync.pdf

General Process Steps:

- Prep your reference image (if needed)
- Create an AutoSync project
- 3. Add your input and reference images
- 4. Add manual control points
- Verify and/or adjust your project parameters
- 6. Solve model
- 7. Run automatic point matching (APM)
- 8. Verify and/or delete control points
- 9. Re-solve model
- 10. Export Image


1. Prep your reference image

You must choose your image wisely

 Ideally this image will be larger than the airphoto(s) you are trying to correct and have similar characteristics.


- For instance, if you are attempting to rectify a color infrared air photo of the Seattle area from 1982, you should find a color infrared reference image (or an image with similar bands) that covers the entire Seattle area. Ideally the spectral signature would be similar if it's a wet year /month you are trying to rectify, pick a reference image taken in a wet year/month.
- You may need to clip out a section of your reference imagery, or mosaic quarter quads together to cover your area of interest. You only get one reference image per project!


2. Create an AutoSync project




3. Add your input and reference images

Your input image is the one you want to rectify, the reference image is the one you want to use as your reference.





In the image above, the upper outside panes have roughly the same size boxes, but different coverage. I will make the right box smaller to compensate:




• Just about perfect!

Now you can adjust the link boxes in the lower panes to about the same size, and use them to zoom in on the image:(look at the inside images on the top) Now we can collect some manual control points to tie the imagery.


4. Add manual control points

- The text box and tools at the bottom of the window is where you do your control point manipulation.
- Make sure you have a wide distribution of points, and a bare minimum of 10 points total (20 is better).
- For the Lamar project, we chose 9 points (if you divided the image into quarters, the corners would be where we tried to put control points) then we added 11 in our area of interest.




5. Verify and/or adjust your project parameters


- Before solving your model or running automatic point matching (APM), you will need to set up your parameters
- Here you will define what strategy you want to use for APM, the geometric correction model, projection, and output settings.



You can read more about modeling in that chapter in the AutoSync manual. For air photos the Direct Linear Transform is typically the best model to use.



- Make sure the radio button for "Same as Reference Image" is selected
- On the "Output" tab, you can make changes to your output and resampling settings - these were defined when you created your project, so only change the settings if you need to.




Last, click back on the first tab, the "APM Strategy" tab.

This tab has a lot of settings, all affecting the generation of automated tie points. You should read the section in the pdf manual "APM Engine" if you have questions.

• Make sure you are matching the correct layer to the correct layer if you have a 3 band image and a 4 band image, pick the bands

that are spectrally closest to one another.


6. Solve model




• Click on the solve button - this will generate the residuals and errors based on your manual tie points. If your RMSE is acceptable, go on to the next step. If not, go through your points and add, delete, or modify as needed.

7. Run automatic point matching (APM)


• Let the fun begin!!!





Oh &*@\$!! What went wrong?

- You have horrible RMSE and none of your points match up what do you do?
- Don't panic!
- Delete the autogenerated points
- Check and adjust your settings
- Try a different band
- Worst case: Start a new project with a different reference image



Advanced Point Mat	ching Strategy		×						
Search Size:	17 💲	Feature Point Density:	100% 💲						
Correlation Size:	11 💲	Minimum Point Match Quality:	0.80						
Least Squares Size:	21 💲	Initial Accuracy:	10%						
✓ Avoid Shadow		Image Scan Type: Positive	e O Negative						
✓ Use Manual Tie Points for Initial Connection between Images									
Exclude Background	d Area	Background Value:	0.00						
Cancel Help									
	Use these se	ttings							

OK, I can live with this RMSE... 8. Verify and/or delete control points


- There are a couple of ways you can go about checking the control points deleting those over a certain error threshold or stepping through your points manually. I like a combination of both.
- Sort your points by error, highest first
- Now you can see what kind of error you are talking about is it just a few outliers, or a bunch of bad matches? Usually this happens because of ground condition changes like fire, phenology, stream changes, landslides, etc.

$\mathbb{R} \setminus \mathbb{C}$) 🚡 🛮 🐯	⊕ ⊕	× ×	⊗ △ '	Σ Σ 🔻		3.00								
		T													-
Point #	Pt. Orig	oint II >	Color	X Input	Y Input	Color	X Ref.	Y Ref.	Z Ref.	X Residual	Y Residual	Error	Contribution	Match	=
1	manual	2 🕨		214.770	898.037		554687.705	4977480.103	1908.664	3.348767	10.987624	11.4860	Column Options	0.000000	
2	manual	3		7957.915	8899.671		568004.098	4963392.708	2277.592	1.045246	0.009233	1.0452		0.000000	
3	manual	4		1675.974	8715.271		557612.828	4964045.473	2736.068	1.781060	-8.214284	8.4051	Select ▶	п попопор	
4	manual	5		9862.007	1073.382		571394.779	4976578.549	2569.001	1.296197	-12.256647	12.3249	Sort •	None þ	
5	manual	6		8888.520	9333.326		569499.298	4962716.543	2412.691	0.870031	-0.524887	1.0161	Edit •	Sort AZ	
6	manual	7		1872.545	9229.413		557885.019	4963161.162	2659.479	-2.308619	-4.552181	5.1041	Format	Sort ZA	
7	manual	8		1270.001	2939.332		556604.123	4973781.515	2012.665	-2.582546	-2.395382	3.5224	Report	Sort ゆ þ	
8	manual	9		4851.624	5601.440		562822.627	4969028.652	2011.018	2.211216	-2.088820	3.0418	Import	0.000000	
9	manual	10		5313.624	5703.858		563614.566	4968838.980	2015.147	-1.882203	-1.320375	2.299	Export	0.000000	
10	manual	11		5858.605	5524.440		564571.543	4969138.267	2064.747	3.799743	-0.610597	3.8484	Formula	0.000000	
11	manual	12		6320.606	5735.256		565369.640	4968765.083	2035.697	5.623317	-3.021097	6.3834	Color	0.000000	
12	manual	13		6557.586	6294.440		565758.835	4967795.789	2045.685	4.503619	-4.264125	6.2020	Alignment •	0.000000	
13	manual	14		6310.887	5549.858		565357.324	4969084.075	2056.895	6.399383	-1.067229	6.4877	Compute Stats	0.000000	
14	manual	15		4588.478	6748.217		562338.596	4967051.884	2028.854	0.910580	-2.920899	3.0595		0.000000	
15	manual	16		5800.295	813.981		564574.007	4977125.393	2511.169	-0.294660	3.316453	3.32951	8 0.000445	0.000000	
16	manual	17		9956.054	5920.654		571232.203	4968425.152	2726.069	-2.894030	-2.952562	4.13437	2 0.000686	0.000000	
17	manual	18		4518.954	5056,459		562254.845	4969984.397	2020.685	-0.116974	-2.014135	2.01752	9 0.000163	0.000000	
18	manual	19		3886.507	5676.197		561145.146	4968915.342	2032.467	-0.109901	2.014895	2.01789	0.000163	0.000000	
19	manual	20		4050.226	4573.527		561451.822	4970831.759	2030.641	-0.330335	0.847902	0.90997	7 0.000033	0.000000	~
<	IIII														>
										RMSE: 8.06	, Error Std. Dev.: 5.	.394 (input) pixels	Cancel		.:

In this
example, there
are some
definite errors.
I would step
through the
points until I
started getting
into "good"
points, then
delete those
points with a
greater error.
Then re-solve
the model.

RinseRepeat

Alternately,
you could step
through each
point and
visually check
it. This
depends on
your time
constraints.




9. Re-solve model


N €	h 🕸	• • × ×	Ø ▲	▼ [Σ] Σ2 [≛	3.00 💲 📕					
Point #	Pt. Orig	Point ID	> Color	X Solve Geome	etric Model using current set c	f points K Ref.	Y Ref.	Z Ref.	X Residual	Y Residual ≙
1	manual	2	•	214.770	898.037	554687.705	4977480.103	1908.664	3.348767	10.98
2	manual	3		7957.915	8899.671	568004.098	4963392.708	2277.592	1.045246	0.00
3	manual	4		1675.974	8715.271	557612.828	4964045.473	2736.068	1.781060	-8.21
4	manual	5		9862.007	1073.382	571394.779	4976578.549	2569.001	1.296197	-12.25
5	manual	6		8888.520	9333.326	569499.298	4962716.543	2412.691	0.870031	-0.52
6	manual	7		1872.545	9229.413	557885.019	4963161.162	2659.479	-2.308619	-4.55
7	manual	8		1270.001	2939.332	556604.123	4973781.515	2012.665	-2.582546	-2.39
8	manual	9		4851.624	5601.440	562822.627	4969028.652	2011.018	2.211216	-2.08
9	manual	10		5313.624	5703.858	563614.566	4968838.980	2015.147	-1.882203	-1.32
10	manual	11		5858.605	5524.440	564571.543	4969138.267	2064.747	3.799743	-0.61
11	manual	12		6320.606	5735.256	565369.640	4968765.083	2035.697	5.623317	-3.02 🕶
										>
RMSE: 7.84, Error Std. Dev.: 5.159 (input) pixels							Cancel		.::	


10. Resample Image (create output image)

It's a good idea to verify your project parameters again - On the "Ŏutput" tab, you can make changes to your output and resampling settings - these were defined when you created your project, so only change the settings if you need to.



Presto!





More Magic...

THE EFFECT OF RIP-RAP ALONG SODA BUTTE CREEK NEAR THE CONFLUENCE OF THE LAMAR RIVER AND AT ROUND PRAIRIE





Figure 1. Photograph of the Lamar River drainage showing grassy river terraces and snowcovered Absaroka Mountains.

By
Cheryl Jaworowski, Dan Mahony, Henry Heasler,
Geology Program,
Yellowstone Center For Resources, Yellowstone National Park
Carrie Guiles
Spatial Analysis Center
Yellowstone Center For Resources, Yellowstone National Park
And
Louis Wasniewski
Caribou-Targhee National Forest

February 22, 2011







Questions?

Carrie Guiles @nps.gov